



FIG. 15

PERCENTAGE OF FGF2 & TGF β NOT DEGRADED
 BY TRYPSIN IN THE PRESENCE OF THE POLYMERS
 DERIVED FROM DEXTRANS

Polymers	% protection of the factors		Polymers	% protection of the factors	
	FGF2	TGF β		FGF2	TGF β
Heparin	100	15	CM ₃ D	22	23
CM ₁ D	20	25	CM ₃ DS _{0.5}	29	32
CM ₁ DS _{0.5}	74	65	CM ₃ DS ₁	32	38
CM ₁ DS _{0.75}	77	71	CM ₃ DS _{1.5}	35	40
CM ₁ DS ₁	80	75	CM ₃ DS ₂	40	47
CM ₁ DS _{1.5}	96	78	CM ₂ DPhSS1	76	67
CM ₁ DS ₂	100	80	CM ₂ DES1	81	71
CM ₁ DSex	100	81	CM ₂ DPhS2	67	56
CM ₂ D	20	25	CM ₃ DTyrS2	83	54
CM ₂ DS _{0.5}	87	74	CM ₁ DPalmS1	67	74
CM ₂ DS _{0.75}	90	77	CM ₁ DOleicS1	58	72
CM ₂ DS ₁	97	80	DS commercial	87	12
CM ₂ DS _{1.5}	95	79	DS _{0.5} equiv	66	9
CM ₂ DS ₂	90	80	DS _{0.125} equiv	51	10
CM ₂ DSex	88	74	Dextran T40	6	5